

MET/HGFR Antibody

Purified Mouse Monoclonal Antibody (Mab)
 Catalog # AM1001a-400 □

Specification

MET/HGFR Antibody - Product info

Application	WB, IF, IHC-P
Primary Accession	P08581
Reactivity	Human, Mouse
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse IgG1
Clone Names	4AT44

MET/HGFR Antibody - Additional info

Gene ID 4233

Other Names

Hepatocyte growth factor receptor, HGF receptor, HGF/SF
 receptor, Proto-oncogene c-Met, Scatter factor receptor, SF
 receptor, Tyrosine-protein kinase Met, MET

Target/Specificity

This monoclonal antibody is generated from mice immunized
 with purified recombinant protein encoding the catalytic
 domain of human Met.

Dilution

WB~~1:8000
 IHC-P~~1:50~100
 IF~~1:100

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V)
 sodium azide. This antibody is purified through a protein G
 column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term
 storage store at -20°C in small aliquots to prevent freeze-thaw
 cycles.

Precautions


MET/HGFR Antibody is for research use only and not for use in
 diagnostic or therapeutic procedures.

MET/HGFR Antibody - Protein Information


Name MET

Function

Receptor tyrosine kinase that transduces signals from the
 extracellular matrix into the cytoplasm by binding to
 hepatocyte growth factor/HGF ligand. Regulates many
 physiological processes including proliferation, scattering,
 morphogenesis and survival. Ligand binding at the cell surface



All lanes : Anti-MET/HGFR Antibody at
 dilution Lane 1: HeLa whole cell lysate
 Lane 2: HepG2 whole cell lysate Lane 3:
 COS-7 whole cell lysate Lysates/proteins
 at 20 µg per lane. Secondary Goat
 Anti-mouse IgG, (H+L), Peroxidase
 conjugated at 1/10000 dilution. Predicted
 band size : 156 kDa Blocking/Dilution
 buffer: 5% NFD/MTBST.



Immunohistochemical analysis of
 paraffin-embedded H.liver section using
 MET/HGFR Antibody(Cat#AM1001a).
 AM1001a was diluted at 1:25 dilution. A
 peroxidase-conjugated goat anti-mouse
 IgG at 1:400 dilution was used as the
 secondary antibody, followed by DAB

induces autophosphorylation of MET on its intracellular domain that provides docking sites for downstream signaling molecules. Following activation by ligand, interacts with the PI3-kinase subunit PIK3R1, PLCG1, SRC, GRB2, STAT3 or the adapter GAB1. Recruitment of these downstream effectors by MET leads to the activation of several signaling cascades including the RAS-ERK, PI3 kinase-AKT, or PLCgamma-PKC. The RAS-ERK activation is associated with the morphogenetic effects while PI3K/AKT coordinates prosurvival effects. During embryonic development, MET signaling plays a role in gastrulation, development and migration of muscles and neuronal precursors, angiogenesis and kidney formation. In adults, participates in wound healing as well as organ regeneration and tissue remodeling. Promotes also differentiation and proliferation of hematopoietic cells. May regulate cortical bone osteogenesis (By similarity).

Cellular Location

Membrane; Single-pass type I membrane protein

Tissue Location

Expressed in normal hepatocytes as well as in epithelial cells lining the stomach, the small and the large intestine. Found also in basal keratinocytes of esophagus and skin. High levels are found in liver, gastrointestinal tract, thyroid and kidney. Also present in the brain. Expressed in metaphyseal bone (at protein level) (PubMed:26637977)

MET/HGFR Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [□ Western Blot](#)
- [□ Blocking Peptides](#)
- [□ Dot Blot](#)
- [□ Immunohistochemistry](#)
- [□ Immunofluorescence](#)
- [□ Immunoprecipitation](#)
- [□ Flow Cytometry](#)
- [□ Cell Culture](#)


MET/HGFR Antibody - Background

The proto-oncogene MET product is the hepatocyte growth factor receptor and encodes tyrosine-kinase activity. The primary single chain precursor protein is post-translationally cleaved to produce the alpha and beta subunits, which are disulfide linked to form the mature receptor. Various mutations in the MET gene are associated with papillary renal carcinoma. Two transcript variants encoding different isoforms have been found for this gene.


MET/HGFR Antibody - References

References for protein: 1. MET receptor sequence variants R970C and T992I lack transforming capacity. Tyner JW, et al. *Cancer Res*, 2010 Aug 1. PMID 20670955. 2. Further evidence for the role of MET in autism susceptibility. Thanseem I, et al. *Neurosci Res*, 2010 Oct. PMID 20615438. 3. Increased HGF and c-Met in muscle tissues of polymyositis and dermatomyositis patients: beneficial roles of HGF in muscle regeneration. Sugiura T, et al. *Clin Immunol*, 2010 Sep. PMID 20580899. 4. Correlation between hepatocyte growth factor receptor and vascular endothelial growth factor-A in breast carcinoma. Gisterek I, et al. *Folia Histochem Cytobiol*, 2010 Jan 1. PMID 20529820. 5. MET overexpressing chordomas frequently exhibit polysomy of chromosome 7 but no MET activation through sarcoma-specific gene fusions. Grabellus F, et al. *Tumour Biol*, 2010 Jun. PMID 20512480. References for HepG2 cell line: 1. Knowles BB,

staining.



Fluorescent confocal image of HepG2 cells stained with MET/HGFR antibody. HepG2 cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.2%, 30 min). Cells were then incubated with AM1001a MET/HGFR primary antibody (1:100, 2 h at room temperature). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-mouse antibody (green) was used (1:1000, 1h). Nuclei were counterstained with Hoechst 33342 (blue) (10 µg/ml, 5 min). Note the highly specific localization of the MET immunosignal to the cytoplasm, supported by Human Protein Atlas Data (<http://www.proteinatlas.org/E NSG00000105976>).



Formalin-fixed and paraffin-embedded human colon carcinoma tissue reacted with MET/HGFR Antibody (Cat.#AM1001a), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

et al. (1980). Human hepatocellular carcinoma cell lines secrete the major plasma proteins and hepatitis B surface antigen. *Science* 209: 497-499.[PubMed: 6248960]. 2. Darlington GJ, et al. (1987). Growth and hepatospecific gene expression of human hepatoma cells in a defined medium. *In Vitro Cell. Dev. Biol.* 23: 349-354.[PubMed: 3034851]. 3. Ihrke, G; Neufeld, EB; Meads, T; Shanks, MR; Cassio, D; Laurent, M; Schroer, TA; Pagano, RE et al. (1993). "WIF-B cells: an in vitro model for studies of hepatocyte polarity". *Journal of Cell Biology* 123 (6): 1761-1775. [PubMed:7506266]. 4. Mersch-Sundermann, V.; Knasmüller, S.; Wu, X. J.; Darroudi, F.; Kassie, F. (2004). "Use of a human-derived liver cell line for the detection of cytoprotective, antigenotoxic and cogenotoxic agents". *Toxicology* 198 (1-3): 329-340. [PubMed:15138059].