

## AKT2 Antibody

Purified Mouse Monoclonal Antibody (Mab)  
 Catalog # AM1848b-400 □

### Specification

#### AKT2 Antibody - Product info

Application	WB, IHC-P, IF
Primary Accession	<a href="#">P31751</a>
Other Accession	<a href="#">NP_001617.1</a>
Reactivity	Human, Mouse, Rat
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse IgG1
Clone Names	148CT5.4.1.4
Calculated MW	55769

#### AKT2 Antibody - Additional info

Gene ID 208

#### Other Names

RAC-beta serine/threonine-protein kinase, Protein kinase Akt-2,  
 Protein kinase B beta, PKB beta, RAC-PK-beta, AKT2

#### Target/Specificity

This AKT2 monoclonal antibody is generated from mouse  
 immunized with AKT2 recombinant protein.

#### Dilution

WB~~1:100~500  
 IHC-P~~1:50~100  
 IF~~1:10~50

#### Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V)  
 sodium azide. This antibody is purified through a protein G  
 column, followed by dialysis against PBS.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term  
 storage store at -20°C in small aliquots to prevent freeze-thaw  
 cycles.

#### Precautions


AKT2 Antibody is for research use only and not for use in  
 diagnostic or therapeutic procedures.

#### AKT2 Antibody - Protein Information


Name AKT2

#### Function

AKT2 is one of 3 closely related serine/threonine- protein  
 kinases (AKT1, AKT2 and AKT3) called the AKT kinase, and which  
 regulate many processes including metabolism, proliferation,  
 cell survival, growth and angiogenesis. This is mediated  
 through serine and/or threonine phosphorylation of a range of



All lanes : Anti-AKT2 Antibody at  
 1:500-1:2000 dilution Lane 1: Jurkat  
 whole cell lysate Lane 2: HeLa whole cell  
 lysate Lane 3: Mouse brain lysate Lane 4:  
 C6 whole cell lysate Lysates/proteins at  
 20 µg per lane. Secondary Goat  
 Anti-mouse IgG, (H+L), Peroxidase  
 conjugated at 1/10000 dilution. Predicted  
 band size : 56 kDa Blocking/Dilution  
 buffer: 5% NFDN/TBST.



AKT2 antibody (Cat. #AM1848b) western  
 blot analysis in mouse brain  
 lysates (15µg/lane).This demonstrates the  
 AKT2 antibody detected the AKT2 protein  
 (arrow).


downstream substrates. Over 100 substrate candidates have been reported so far, but for most of them, no isoform specificity has been reported. AKT is responsible of the regulation of glucose uptake by mediating insulin-induced translocation of the SLC2A4/GLUT4 glucose transporter to the cell surface. Phosphorylation of PTPN1 at 'Ser-50' negatively modulates its phosphatase activity preventing dephosphorylation of the insulin receptor and the attenuation of insulin signaling. Phosphorylation of TBC1D4 triggers the binding of this effector to inhibitory 14-3-3 proteins, which is required for insulin-stimulated glucose transport. AKT regulates also the storage of glucose in the form of glycogen by phosphorylating GSK3A at 'Ser-21' and GSK3B at 'Ser-9', resulting in inhibition of its kinase activity. Phosphorylation of GSK3 isoforms by AKT is also thought to be one mechanism by which cell proliferation is driven. AKT regulates also cell survival via the phosphorylation of MAP3K5 (apoptosis signal-related kinase). Phosphorylation of 'Ser-83' decreases MAP3K5 kinase activity stimulated by oxidative stress and thereby prevents apoptosis. AKT mediates insulin-stimulated protein synthesis by phosphorylating TSC2 at 'Ser-939' and 'Thr-1462', thereby activating mTORC1 signaling and leading to both phosphorylation of 4E-BP1 and in activation of RPS6KB1. AKT is involved in the phosphorylation of members of the FOXO factors (Forkhead family of transcription factors), leading to binding of 14-3-3 proteins and cytoplasmic localization. In particular, FOXO1 is phosphorylated at 'Thr-24', 'Ser-256' and 'Ser-319'. FOXO3 and FOXO4 are phosphorylated on equivalent sites. AKT has an important role in the regulation of NF-kappa-B-dependent gene transcription and positively regulates the activity of CREB1 (cyclic AMP (cAMP)- response element binding protein). The phosphorylation of CREB1 induces the binding of accessory proteins that are necessary for the transcription of pro-survival genes such as BCL2 and MCL1. AKT phosphorylates 'Ser-454' on ATP citrate lyase (ACLY), thereby potentially regulating ACLY activity and fatty acid synthesis. Activates the 3B isoform of cyclic nucleotide phosphodiesterase (PDE3B) via phosphorylation of 'Ser-273', resulting in reduced cyclic AMP levels and inhibition of lipolysis. Phosphorylates PIKFYVE on 'Ser-318', which results in increased PI(3)P-5 activity. The Rho GTPase-activating protein DLC1 is another substrate and its phosphorylation is implicated in the regulation cell proliferation and cell growth. AKT plays a role as key modulator of the AKT-mTOR signaling pathway controlling the tempo of the process of newborn neurons integration during adult neurogenesis, including correct neuron positioning, dendritic development and synapse formation. Signals downstream of phosphatidylinositol 3-kinase (PI(3)K) to mediate the effects of various growth factors such as platelet-derived growth factor (PDGF), epidermal growth factor (EGF), insulin and insulin-like growth factor I (IGF-I). AKT mediates the antiapoptotic effects of IGF-I. Essential for the SPATA13-mediated regulation of cell migration and adhesion assembly and disassembly. May be involved in the regulation of the placental development.

#### Cellular Location


Cytoplasm. Nucleus. Cell membrane; Peripheral membrane protein. Early endosome {ECO:0000250|UniProtKB:Q60823}. Note=Localizes within both nucleus and cytoplasm of proliferative primary myoblasts and mostly within the nucleus of differentiated primary myoblasts. By virtue of the N-terminal PH domain, is recruited to sites of the plasma membrane containing increased PI(3,4,5)P3 or PI(3,4)P2, cell membrane targeting is also facilitated by interaction with CLIP3 Colocalizes with WDFY2 in early endosomes (By similarity) {ECO:0000250|UniProtKB:Q60823}

#### Tissue Location

Expressed in all cell types so far analyzed.



AKT2 Monoclonal Antibody (Cat. #AM1848b) immunohistochemistry analysis in formalin fixed and paraffin embedded human prostate carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the AKT2 Monoclonal Antibody for immunohistochemistry. Clinical relevance has not been evaluated.



Confocal immunofluorescent analysis of AKT2 Antibody (Cat#AM1848b) with HeLa cell followed by Alexa Fluor® 488-conjugated goat anti-mouse IgG (green). DAPI was used to stain the cell nuclear (blue).

## AKT2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [□Western Blot](#)
- [□Blocking Peptides](#)
- [□Dot Blot](#)
- [□Immunohistochemistry](#)
- [□Immunofluorescence](#)
- [□Immunoprecipitation](#)
- [□Flow Cytometry](#)
- [□Cell Culture](#)

## AKT2 Antibody - Background

AKT2 is a putative oncogene encoding a protein belonging to a subfamily of serine/threonine kinases containing SH2-like (Src homology 2-like) domains. AKT2 was shown to be amplified and overexpressed in 2 of 8 ovarian carcinoma cell lines and 2 of 15 primary ovarian tumors. Overexpression contributes to the malignant phenotype of a subset of human ductal pancreatic cancers. The encoded protein is a general protein kinase capable of phosphorylating several known proteins.

## AKT2 Antibody - References

Fortier, A.M., et al. FEBS Lett. 584(5):984-988(2010) Li, X., et al. Arterioscler. Thromb. Vasc. Biol. 30(3):582-590(2010) Santi, S.A., et al. Am. J. Physiol., Cell Physiol. 298 (3), C580-C591 (2010)

## AKT2 Antibody - Citations

- [AKT2-knockdown suppressed viability with enhanced apoptosis, and attenuated chemoresistance to temozolomide of human glioblastoma cells in vitro and in vivo.](#)
- [MiRNA-615-5p Functions as a Tumor Suppressor in Pancreatic Ductal Adenocarcinoma by Targeting AKT2.](#)