

## Phospho-MET(Y1356) Antibody

Peptide Affinity Purified Rabbit Polyclonal Antibody (Pab)  
Catalog # AP3168a-400 □

### Specification

#### Phospho-MET(Y1356) Antibody - Product info

Application	DB
Primary Accession	<a href="#">P08581</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Clone Names	RB24512
Calculated MW	155541

#### Phospho-MET(Y1356) Antibody - Additional info

Gene ID 4233

#### Other Names

Hepatocyte growth factor receptor, HGF receptor, HGF/SF receptor, Proto-oncogene c-Met, Scatter factor receptor, SF receptor, Tyrosine-protein kinase Met, MET

#### Target/Specificity

This MET Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding Y1356 of human MET.

#### Dilution

DB~1:500

#### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions


Phospho-MET(Y1356) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

#### Phospho-MET(Y1356) Antibody - Protein Information

Name MET

#### Function

Receptor tyrosine kinase that transduces signals from the extracellular matrix into the cytoplasm by binding to hepatocyte growth factor/HGF ligand. Regulates many physiological processes including proliferation, scattering, morphogenesis and survival. Ligand binding at the cell surface induces autophosphorylation of MET on its intracellular



domain that provides docking sites for downstream signaling molecules. Following activation by ligand, interacts with the PI3-kinase subunit PIK3R1, PLCG1, SRC, GRB2, STAT3 or the adapter GAB1. Recruitment of these downstream effectors by MET leads to the activation of several signaling cascades including the RAS-ERK, PI3 kinase-AKT, or PLCgamma-PKC. The RAS-ERK activation is associated with the morphogenetic effects while PI3K/AKT coordinates prosurvival effects. During embryonic development, MET signaling plays a role in gastrulation, development and migration of muscles and neuronal precursors, angiogenesis and kidney formation. In adults, participates in wound healing as well as organ regeneration and tissue remodeling. Promotes also differentiation and proliferation of hematopoietic cells. May regulate cortical bone osteogenesis (By similarity).

#### Cellular Location

Membrane; Single-pass type I membrane protein

#### Tissue Location

Expressed in normal hepatocytes as well as in epithelial cells lining the stomach, the small and the large intestine. Found also in basal keratinocytes of esophagus and skin. High levels are found in liver, gastrointestinal tract, thyroid and kidney. Also present in the brain. Expressed in metaphyseal bone (at protein level) (PubMed:26637977)

### Phospho-MET(Y1356) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [□Western Blot](#)
- [□Blocking Peptides](#)
- [□Dot Blot](#)
- [□Immunohistochemistry](#)
- [□Immunofluorescence](#)
- [□Immunoprecipitation](#)
- [□Flow Cytometry](#)
- [□Cell Culture](#)

### Phospho-MET(Y1356) Antibody - Background

The proto-oncogene MET product is the hepatocyte growth factor receptor and encodes tyrosine-kinase activity. The primary single chain precursor protein is post-translationally cleaved to produce the alpha and beta subunits, which are disulfide linked to form the mature receptor. Various mutations in the MET gene are associated with papillary renal carcinoma.

### Phospho-MET(Y1356) Antibody - References

Wong, A.S., et al., Exp. Cell Res. 299(1):248-256 (2004). Higuchi, T., et al., Mol. Cell. Biol. 24(17):7456-7468 (2004). Mineo, R., et al., Endocrinology 145(9):4355-4365 (2004). Chung, J., et al., J. Biol. Chem. 279(31):32287-32293 (2004). Fischer, O.M., et al., J. Biol. Chem. 279(28):28970-28978 (2004).

#### Phospho-MET(Y1356) Antibody - Citations

- [Receptor-type protein tyrosine phosphatase beta \(RPTP-beta\) directly dephosphorylates and regulates hepatocyte growth factor receptor \(HGFR/Met\) function.](#)