

## AKAP8 Antibody

Purified Rabbit Polyclonal Antibody (Pab)  
Catalog # AP51662-100 □

### Specification

#### AKAP8 Antibody - Product info

Application	WB
Primary Accession	<a href="#">Q43823</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	95 KDa

#### AKAP8 Antibody - Additional info

Gene ID 10270

#### Other Names

A-kinase anchor protein 8, AKAP-8, A-kinase anchor protein 95 kDa, AKAP 95, AKAP8, AKAP95

#### Target/Specificity

KLH conjugated synthetic peptide derived from human AKAP8

#### Dilution

WB~ 1:1000

#### Format

0.01M PBS, pH 7.2, 0.1% Sodium azide, Glycerol 50%

#### Storage

Store at -20 °C. Stable for 12 months from date of receipt


#### AKAP8 Antibody - Protein Information

Name AKAP8

Synonyms AKAP95

#### Function

Anchoring protein that mediates the subcellular compartmentation of cAMP-dependent protein kinase (PKA type II) (PubMed:<a href="http://www.uniprot.org/citations/9473338" target="\_blank">9473338</a>). Acts as an anchor for a PKA-signaling complex onto mitotic chromosomes, which is required for maintenance of chromosomes in a condensed form throughout mitosis. Recruits condensin complex subunit NCAPD2 to chromosomes required for chromatin condensation; the function appears to be independent from PKA-anchoring (PubMed:<a href="http://www.uniprot.org/citations/10601332" target="\_blank">10601332</a>, PubMed:<a href="http://www.uniprot.org/citations/10791967" target="\_blank">10791967</a>, PubMed:<a href="http://www.uniprot.org/citations/11964380" target="\_blank">11964380</a>). May help to deliver cyclin D/E



to CDK4 to facilitate cell cycle progression (PubMed:<a href="http://www.uniprot.org/citations/14641107" target="\_blank">14641107</a>). Required for cell cycle G2/M transition and histone deacetylation during mitosis. In mitotic cells recruits HDAC3 to the vicinity of chromatin leading to deacetylation and subsequent phosphorylation at 'Ser-10' of histone H3; in this function may act redundantly with AKAP8L (PubMed:<a href="http://www.uniprot.org/citations/16980585" target="\_blank">16980585</a>). Involved in nuclear retention of RPS6KA1 upon ERK activation thus inducing cell proliferation (PubMed:<a href="http://www.uniprot.org/citations/22130794" target="\_blank">22130794</a>). May be involved in regulation of DNA replication by acting as scaffold for MCM2 (PubMed:<a href="http://www.uniprot.org/citations/12740381" target="\_blank">12740381</a>). Enhances HMT activity of the KMT2 family MLL4/WBP7 complex and is involved in transcriptional regulation. In a teratocarcinoma cell line is involved in retinoic acid-mediated induction of developmental genes implicating H3 'Lys-4' methylation (PubMed:<a href="http://www.uniprot.org/citations/23995757" target="\_blank">23995757</a>). May be involved in recruitment of active CASP3 to the nucleus in apoptotic cells (PubMed:<a href="http://www.uniprot.org/citations/16227597" target="\_blank">16227597</a>). May act as a carrier protein of GJA1 for its transport to the nucleus (PubMed:<a href="http://www.uniprot.org/citations/26880274" target="\_blank">26880274</a>). Seems to be involved in modulation of rDNA transcription. Preferentially binds GC-rich DNA in vitro and associates to GC-rich ribosomal RNA promoters (PubMed:<a href="http://www.uniprot.org/citations/26683827" target="\_blank">26683827</a>). Involved in modulation of Toll-like receptor signaling. Required for the cAMP-dependent suppression of TNF-alpha in early stages of LPS-induced macrophage activation; the function probably implicates targeting of PKA to NFkB1 (By similarity).

#### Cellular Location

Nucleus. Nucleus matrix. Nucleus, nucleolus. Cytoplasm {ECO:0000250|UniProtKB:Q9DBR0}. Note=Associated with the nuclear matrix in interphase and redistributes mostly to chromatin at mitosis. However, mitotic chromatin localization has been questioned. Upon nuclear reassembly at the end of mitosis, is sequestered into the daughter nuclei where it re-acquires an interphase distribution. Localized to the nucleolus in interphase Colocalizes with GJA1 at the nuclear membrane specifically during cell cycle G1/S phase.

#### Tissue Location

Highly expressed in heart, liver, skeletal muscle, kidney and pancreas. Expressed in mature dendritic cells

## AKAP8 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [□Western Blot](#)
- [□Blocking Peptides](#)
- [□Dot Blot](#)
- [□Immunohistochemistry](#)
- [□Immunofluorescence](#)
- [□Immunoprecipitation](#)
- [□Flow Cytometry](#)
- [□Cell Culture](#)

## AKAP8 Antibody - Background

Anchoring protein that mediates the subcellular

compartmentation of cAMP-dependent protein kinase (PKA type II).

#### AKAP8 Antibody - References

Eide T.,et al.Exp. Cell Res. 238:305-316(1998). Grimwood J.,et al.Nature 428:529-535(2004). Olsen J.V.,et al.Cell 127:635-648(2006). Daub H.,et al.Mol. Cell 31:438-448(2008). Dephoure N.,et al.Proc. Natl. Acad. Sci. U.S.A. 105:10762-10767(2008).