

Anti-Met Antibody

Mouse Monoclonal Antibody
Catalog # AP53480-100 □

Specification

Anti-Met Antibody - Product Information

Application	WB
Primary Accession	P08581
Other Accession	NM_000245
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2a
Immunogen	Purified recombinant human Met protein expressed in E.coli.
Purification	Affinity purified
Calculated MW	156KDa KDa

Anti-Met Antibody - Additional Information

Gene ID 4233

Other Names

AUTS9;c met;D249;Hepatocyte growth factor receptor;HGF;HGF receptor;HGF/SF receptor;HGFR;MET;Met proto oncogene tyrosine kinase;MET proto oncogene, receptor tyrosine kinase;Met proto-oncogene (hepatocyte growth factor receptor);Met proto-oncogene;Met protooncogene;MET_HUMAN;Oncogene MET;Par4;Proto-oncogene c-Met;RCCP2;Scatter factor receptor;SF receptor;Tyrosine-protein kinase Met.

Dilution

WB~1:1000

Format

Purified mouse monoclonal antibody in PBS(pH 7.4) containing with 0.03% Proclin300 and 50% glycerol.

Storage


Store at -20 °C.Stable for 12 months from date of receipt

Anti-Met Antibody - Protein Information

Name MET

Function

Receptor tyrosine kinase that transduces signals from the extracellular matrix into the cytoplasm by binding to hepatocyte growth factor/HGF ligand. Regulates many physiological processes including proliferation, scattering, morphogenesis and survival. Ligand binding at the cell surface induces autophosphorylation of MET on its intracellular domain that provides docking sites for downstream signaling molecules. Following activation by ligand, interacts with the PI3-kinase subunit PIK3R1, PLCG1, SRC, GRB2, STAT3 or the adapter GAB1. Recruitment of these downstream effectors by



MET leads to the activation of several signaling cascades including the RAS-ERK, PI3 kinase-AKT, or PLCgamma-PKC. The RAS-ERK activation is associated with the morphogenetic effects while PI3K/AKT coordinates prosurvival effects. During embryonic development, MET signaling plays a role in gastrulation, development and migration of muscles and neuronal precursors, angiogenesis and kidney formation. In adults, participates in wound healing as well as organ regeneration and tissue remodeling. Promotes also differentiation and proliferation of hematopoietic cells. May regulate cortical bone osteogenesis (By similarity).

Cellular Location

Membrane; Single-pass type I membrane protein

Tissue Location

Expressed in normal hepatocytes as well as in epithelial cells lining the stomach, the small and the large intestine. Found also in basal keratinocytes of esophagus and skin. High levels are found in liver, gastrointestinal tract, thyroid and kidney. Also present in the brain. Expressed in metaphyseal bone (at protein level) (PubMed:26637977)

Anti-Met Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [□ Western Blot](#)
- [□ Blocking Peptides](#)
- [□ Dot Blot](#)
- [□ Immunohistochemistry](#)
- [□ Immunofluorescence](#)
- [□ Immunoprecipitation](#)
- [□ Flow Cytometry](#)
- [□ Cell Culture](#)

Anti-Met Antibody - Background

Receptor for hepatocyte growth factor and scatter factor. Has a tyrosine-protein kinase activity. Functions in cell proliferation, scattering, morphogenesis and survival.