

CEA Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6513c-400 □

Specification

CEA Antibody (Center) - Product info

Application	IHC-P-Leica, FC, WB
Primary Accession	P06731
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Clone Names	RB19418
Calculated MW	76795

CEA Antibody (Center) - Additional info

Gene ID 1048

Other Names

Carcinoembryonic antigen-related cell adhesion molecule 5,
Carcinoembryonic antigen, CEA, Meconium antigen 100,
CD66e, CEACAM5, CEA

Target/Specificity

This CEA antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 400-429 amino acids from the Central region of human CEA.

Dilution

IHC-P-Leica~~1:500
WB~~1:1000
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CEA Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.


CEA Antibody (Center) - Protein Information

Name CEACAM5


Synonyms CEA

Function

Cell surface glycoprotein that plays a role in cell adhesion, intracellular signaling and tumor progression (PubMed:<a



Immunohistochemical analysis of paraffin-embedded Human colon carcinoma tissue using AP6513C performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9.0). Samples were incubated with primary antibody(1:500) for 1 hours at room temperature. A undiluted biotinylated CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



All lanes : Anti-CEA Antibody (Center) at 1:1000 dilution Lane 1: MCF-7 whole cell lysate Lane 2: HT-29 whole cell lysate

<http://www.uniprot.org/citations/2803308>, PubMed:10910050, PubMed:10864933). Mediates homophilic and heterophilic cell adhesion with other carcinoembryonic antigen-related cell adhesion molecules, such as CEACAM6 (PubMed:2803308). Plays a role as an oncogene by promoting tumor progression; induces resistance to anoikis of colorectal carcinoma cells (PubMed:10910050).

Cellular Location

Cell membrane; Lipid- anchor, GPI-anchor. Apical cell membrane. Cell surface Note=Localized to the apical glycocalyx surface

Tissue Location

Expressed in columnar epithelial and goblet cells of the colon (at protein level) (PubMed:10436421). Found in adenocarcinomas of endoderally derived digestive system epithelium and fetal colon.

CEA Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [□Western Blot](#)
- [□Blocking Peptides](#)
- [□Dot Blot](#)
- [□Immunohistochemistry](#)
- [□Immunofluorescence](#)
- [□Immunoprecipitation](#)
- [□Flow Cytometry](#)
- [□Cell Culture](#)


CEA Antibody (Center) - Background

CEA is a cell surface glycoprotein that plays a role in cell adhesion and in intracellular signaling. It is a receptor for E.coli Dr adhesins.

CEA Antibody (Center) - References

Lagoudianakis,E.E., J. Surg. Res. 155 (2), 183-190 (2009)
Gambichler,T., Am. J. Clin. Pathol. 131 (6), 782-787 (2009)

Lane 3: PC-3 whole cell lysate Lane 4: A549 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 77 kDa Blocking/Dilution buffer: 5% NFDN/TBST.



CEA Antibody (Center) (Cat. #AP6513c) flow cytometric analysis of 293 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.