

UHRF1 Antibody (Center)

Peptide Affinity Purified Rabbit Polyclonal Antibody (Pab)
 Catalog # AP8846c-400 □

Specification

UHRF1 Antibody (Center) - Product info

Application	FC, IF, WB
Primary Accession	Q96T88
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	89814

UHRF1 Antibody (Center) - Additional info

Gene ID 29128

Other Names

E3 ubiquitin-protein ligase UHRF1, 632-, Inverted CCAAT box-binding protein of 90 kDa, Nuclear protein 95, Nuclear zinc finger protein Np95, HuNp95, hNp95, RING finger protein 106, Transcription factor ICBP90, Ubiquitin-like PHD and RING finger domain-containing protein 1, hUHRF1, Ubiquitin-like-containing PHD and RING finger domains protein 1, UHRF1, ICBP90, NP95, RNF106

Target/Specificity

This UHRF1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 229-257 amino acids from the Central region of human UHRF1.

Dilution

WB~~1:1000

FC~~1:10~50

IF~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.


Precautions

UHRF1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.


UHRF1 Antibody (Center) - Protein Information

Name UHRF1

Synonyms ICBP90, NP95, RNF106



UHRF1 Antibody (Center) (Cat. #AP8846c) western blot analysis in HeLa cell line lysates (35ug/lane). This demonstrates the UHRF1 antibody detected the UHRF1 protein (arrow).



UHRF1 Antibody (Center) (Cat.#AP8846c) flow cytometry analysis of HeLa cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Function

Multidomain protein that acts as a key epigenetic regulator by bridging DNA methylation and chromatin modification. Specifically recognizes and binds hemimethylated DNA at replication forks via its YDG domain and recruits DNMT1 methyltransferase to ensure faithful propagation of the DNA methylation patterns through DNA replication. In addition to its role in maintenance of DNA methylation, also plays a key role in chromatin modification: through its tudor-like regions and PHD- type zinc fingers, specifically recognizes and binds histone H3 trimethylated at 'Lys-9' (H3K9me3) and unmethylated at 'Arg-2' (H3R2me0), respectively, and recruits chromatin proteins. Enriched in pericentric heterochromatin where it recruits different chromatin modifiers required for this chromatin replication. Also localizes to euchromatic regions where it negatively regulates transcription possibly by impacting DNA methylation and histone modifications. Has E3 ubiquitin-protein ligase activity by mediating the ubiquitination of target proteins such as histone H3 and PML. It is still unclear how E3 ubiquitin-protein ligase activity is related to its role in chromatin in vivo. May be involved in DNA repair.

Cellular Location

Nucleus {ECO:0000255 | PROSITE- ProRule:PRU00358, ECO:0000269 | PubMed:10646863, ECO:0000269 | PubMed:17673620, ECO:0000269 | PubMed:17967883, ECO:0000269 | PubMed:19056828, ECO:0000269 | PubMed:21777816} Note=Localizes to replication foci. Enriched in pericentric heterochromatin. Also localizes to euchromatic regions

Tissue Location

Expressed in thymus, bone marrow, testis, lung and heart.
Overexpressed in breast cancer



Fluorescent confocal image of HeLa cell stained with UHRF1 Antibody (Center)(Cat#AP8846c). HeLa cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.1%, 10 min), then incubated with UHRF1 primary antibody (1:25, 1 h at 37°C). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:400, 50 min at 37°C). Cytoplasmic actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (7 units/ml, 1 h at 37°C). Nuclei were counterstained with DAPI (blue) (10 µg/ml, 10 min). UHRF1 immunoreactivity is localized to Nucleus significantly.

UHRF1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [□ Western Blot](#)
- [□ Blocking Peptides](#)
- [□ Dot Blot](#)
- [□ Immunohistochemistry](#)
- [□ Immunofluorescence](#)
- [□ Immunoprecipitation](#)
- [□ Flow Cytometry](#)
- [□ Cell Culture](#)

UHRF1 Antibody (Center) - Background

UHRF1 is a member of a subfamily of RING-finger type E3 ubiquitin ligases. The protein binds to specific DNA sequences, and recruits a histone acetylase to regulate gene expression. Its expression peaks at late G1 phase and continues during G2 and M phases of the cell cycle. It plays a major role in the G1/S transition by regulating topoisomerase IIalpha and retinoblastoma gene expression, and functions in the p53-dependent DNA damage checkpoint.

UHRF1 Antibody (Center) - References

Achour, M., et al., Biochem. Biophys. Res. Commun. 390 (3), 523-528 (2009)