

PSMA5 Antibody

Mouse Monoclonal Antibody (Mab)
 Catalog # AW5056-400 □

Specification

PSMA5 Antibody - Product Information

Application	IF, IHC-P, WB
Primary Accession	P28066
Other Accession	Q9Z2U1 , Q5E987 , NP_002781.2
Reactivity	Human
Predicted	Mouse, Rat, Bovine
Host	Mouse
Clonality	Monoclonal
Calculated MW	H=26,20;M=26;Rat=26 KDa
Isotype	IgG1
Antigen Source	HUMAN

PSMA5 Antibody - Additional Information

Gene ID 5686

Antigen Region
 1-261

Other Names
 PSMA5; Proteasome subunit alpha type-5; Macropain zeta chain; Multicatalytic endopeptidase complex zeta chain; Proteasome zeta chain

Dilution
 WB~~1:1000
 IHC-P~~1:25
 IF~~1:25

Target/Specificity
 Purified His-tagged PSMA5 protein(Fragment) was used to produced this monoclonal antibody.


Format
 Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage
 Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.


Precautions
 PSMA5 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

PSMA5 Antibody - Protein Information

Name PSMA5



Western blot analysis of lysates from human liver tissue, MCF-7, HepG2, HeLa, rat PC-12 cell line (from left to right), using PSMA5 Antibody (Cat. #AW5056). AW5056 was diluted at 1:1000 at each lane. A goat anti-mouse IgG H&L (HRP) at 1:10000 dilution was used as the secondary antibody.



Immunohistochemical analysis of paraffin-embedded H. testis section using PSMA5 Antibody (Cat#AW5056). AW5056 was diluted at 1:25 dilution. A peroxidase-conjugated goat anti-mouse IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.

Function

Component of the 20S core proteasome complex involved in the proteolytic degradation of most intracellular proteins. This complex plays numerous essential roles within the cell by associating with different regulatory particles. Associated with two 19S regulatory particles, forms the 26S proteasome and thus participates in the ATP-dependent degradation of ubiquitinated proteins. The 26S proteasome plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins that could impair cellular functions, and by removing proteins whose functions are no longer required. Associated with the PA200 or PA28, the 20S proteasome mediates ubiquitin-independent protein degradation. This type of proteolysis is required in several pathways including spermatogenesis (20S-PA200 complex) or generation of a subset of MHC class I-presented antigenic peptides (20S-PA28 complex).

Cellular Location

Cytoplasm. Nucleus

Tissue Location

Expressed in fetal brain (at protein level).

PSMA5 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [□ Western Blot](#)
- [□ Blocking Peptides](#)
- [□ Dot Blot](#)
- [□ Immunohistochemistry](#)
- [□ Immunofluorescence](#)
- [□ Immunoprecipitation](#)
- [□ Flow Cytometry](#)
- [□ Cell Culture](#)

PSMA5 Antibody - Background

The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit.

PSMA5 Antibody - References

Kottgen, A., et al. *Nat. Genet.* 42(5):376-384(2010) Sugiyama, N., et al. *Mol. Cell Proteomics* 6(6):1103-1109(2007) Olsen, J.V., et al. *Cell* 127(3):635-648(2006) Beausoleil, S.A., et al. *Nat. Biotechnol.* 24(10):1285-1292(2006) Hirano, Y., et al. *Nature* 437(7063):1381-1385(2005)



Fluorescent image of HeLa cells stained with XAF1 PSMA5 Antibody (Cat#AW5056). AW5056 was diluted at 1:25 dilution. An Alexa Fluor 488-conjugated goat anti-mouse IgG at 1:400 dilution was used as the secondary antibody (green). Cytoplasmic actin was counterstained with Alexa Fluor® 555 conjugated with Phalloidin (red).