

## CRABP2 Antibody (C-term)

Peptide Affinity Purified Rabbit Polyclonal Antibody (Pab)  
Catalog # AW5524-200 □

### Specification

#### CRABP2 Antibody (C-term) - Product Information

Application	WB
Primary Accession	<a href="#">P29373</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=16;M=16;R=16 KDa
Isotype	Rabbit Ig
Antigen Source	HUMAN

#### CRABP2 Antibody (C-term) - Additional Information

Gene ID 1382

Antigen Region  
102-136

#### Other Names

Cellular retinoic acid-binding protein 2, Cellular retinoic acid-binding protein II, CRABP-II, CRABP2

#### Dilution

WB~1:1000

#### Target/Specificity

This CRABP2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 102-136 amino acids from the C-terminal region of human CRABP2.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

CRABP2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

#### CRABP2 Antibody (C-term) - Protein Information


Name CRABP2

#### Function

Transports retinoic acid to the nucleus. Regulates the access of retinoic acid to the nuclear retinoic acid receptors.

#### Cellular Location

Cytoplasm. Endoplasmic reticulum. Nucleus. Note=Upon ligand binding, a conformation change exposes a nuclear localization motif and the protein is transported into the nucleus



## CRABP2 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [□Western Blot](#)
- [□Blocking Peptides](#)
- [□Dot Blot](#)
- [□Immunohistochemistry](#)
- [□Immunofluorescence](#)
- [□Immunoprecipitation](#)
- [□Flow Cytometry](#)
- [□Cell Culture](#)

## CRABP2 Antibody (C-term) - Background

A number of specific carrier proteins for members of the vitamin A family have been discovered. Cellular retinoic acid binding proteins (CRABP) are low molecular weight proteins whose precise function remains unknown. The inducibility of the CRABP2 gene suggests that this isoform is important in retinoic acid-mediated regulation of human skin growth and differentiation. It has been postulated that the CRABP2 gene is transcriptionally regulated by a newly synthesized regulatory protein. [provided by RefSeq].

## CRABP2 Antibody (C-term) - References

Sola, R., et al. *Atherosclerosis* 211(2):630-637(2010) Manolescu, D.C., et al. *Pediatr. Res.* 67(6):598-602(2010) Calmon, M.F., et al. *Neoplasia* 11(12):1329-1339(2009) Corlazzoli, F., et al. *PLoS ONE* 4(1), E4305 (2009) : Gupta, A., et al. *Exp. Cell Res.* 314(20):3663-3668(2008)