

HIS Tag Antibody

Mouse Monoclonal Antibody (Mab)
 Catalog # AW5672-200 □

Specification

HIS Tag Antibody - Product Information

Application	WB
Host	Mouse
Clonality	Monoclonal
Calculated MW	45-50KD KDa
Isotype	Mouse IgG1
Antigen Source	HUMAN

HIS Tag Antibody - Additional Information

Antigen Region
 NA

Dilution
 WB~~1:8000

Target/Specificity

Purified recombinant HIS-tagged fusion protein and poly-HIS peptide were used to produced this monoclonal antibody.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HIS Tag Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

HIS Tag Antibody - Protein Information


HIS Tag Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HIS Tag Antibody - Background

Epitope tags consisting of short sequences recognized by well-characterized monoclonal antibodies have been widely used in the study of protein expression in various systems. The 6xHIS tag



All lanes : Anti-HIS Tag at 1:1000 dilution (upper) or GAPDH (lower) Lane 1: 293T/17 transfected with 12tag lysate (10ug) Lane 2: Non-transfected 293T/17 lysate (10ug) Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 45-50 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



Anti-HIS Tag at 1:8000 dilution + 12-tag lysate Lysates/proteins at 20ng per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 48 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

(HHHHHH), recognized by the monoclonal antibody clone 6AT18 provides an established example of this application. 6xHIS-tagged fusion proteins are easily purified from cell lysates by affinity chromatography using Nickel-Sepharose resin. Abgent's anti-6xHIS monoclonal antibody provides a simple solution to detect the expression of HIS-tagged fusion proteins in cells.

HIS Tag Antibody - References

Hochuli E, Doebeli H, and Schacher A. New metal chelate absorbent selective for proteins and peptides containing neighboring histidine residues. *J. Chromatogr.* 1987;411:177-184.